

An annotated checklist of aquatic avifauna of Rajura, Godada and Dhanora lakes of Buldhana district of (M. S.) India

Prasanna Subhash Joshi

Department of Zoology
Dr. R. B. Shingane Science College, Jalgaon (Jamod), Dist- Buldhana (M.S.)
Psjoshi009@gmail.com

ABSTRACT

Abundance of avifauna indicates the healthy status of lake. Observations were made on the occurrence, abundance, richness of avifauna in three lakes viz. Rajura, Godada and Dhanora lakes of Buldhana district in the Indian state Maharashtra. Surveys were conducted during 15 October to 15 November (2011). During study total twenty-eight species of water birds of thirteen families were recorded. The occurrence, ecological status and food habit of different species of aquatic birds was recorded. The studies on aquatic avifauna of Survey district revealed abundance of twenty-eight species of water birds of thirteen families (Table-1). Of all the families, Podicipitidae (1 Species), Phalacrocorax (1 species), Ardeidae (5 species), Ciconiidae (3 species), Treskornithidae (1 species), Anatidae (1 species), Rallidae (3 species), Recurvirostridae (1 species), Charadriidae (2 species) Scolopacidae (3 species), Laridae (1 species), Acledinidae (2 species) and Motacillidae (3 species). It is expected that this study would provide a preliminary database for the aquatic avifauna of this area for the further research.

Key words- Avifauna, Abundance, Occurrence, Water birds.

INTRODUCTION

Nearly everyone enjoys birds: the beauty of their forms and coloring, the vivacity of their movement, the buoyancy of their flight and sweetness of their songs. Birds are often common denizens of the ecosystems and they have been considered as an indicator species of inhabited areas (Blair, 1999). Population of birds is a sensitive indicator of pollution in both terrestrial and aquatic ecosystem (Gaston, 1975; Hardy *et al*, 1987). Wetlands are the most productive and biologically diverse in the world but very fragile ecosystems (Gibbs, 1993). Wetlands and waterbirds are inseparable elements and support a rich array of waterbird communities (Grimmett and Inskipp, 2007). Waterbirds are an important component of most of the wetland ecosystem as they occupy several trophic levels in the food web of wetland nutrient cycles. Activities of water birds are considered as indicator of quality of the wetland ecosystem and form the terminal links in many aquatic food chains, and as a result they reflect changes originating in several different ecosystem components (Custer and Osborne 1977). The various lakes and wetlands in any city serve as a balancing reservoir for sustaining native flora and fauna (Grimmett and Inskipp, 2007; Surana *et al*, 2007). The aquatic birds of the Jalgaon (Jamod) are important bioindicators of lake ecosystems which should be protected to conserve the biodiversity and environment.

MATERIALS AND METHODS

The observations were done at Rajura, Godada and Dhanora lakes of Buldhana district in the Indian state Maharashtra. 'Rajura Lake' lies between latitudinal parallels 21°04'29"N and longitudinal parallels 76°29'31"E at an elevation of 291 m above sea level. The 'Rajura Reservoir' covers area around 5.90 Sq. Km. Dhanora (Hattipaul) Lake located at distance of 10 km. from Jalgaon (Jamod) Tahsil and 55 km. from Khamgaon, District Buldhana (MS.) India. The lake lies between 21° 07' 00" N latitude and 76° 27' 45" E longitude. Lake covers 31.20 hectares surface area with 268 m. length. Godada Lake' lies between latitudinal parallels 21° 05' 45" N and longitudinal parallels 76°31'00" E Km. Lake covers area around 34.64 hectares. During study duration temperature was ranges between 35 to 16°C. An efficient protocol has been adopted (Turner, 2003). Bird sampling was made by walking at a slow pace (about 1-1.5 km hr⁻¹) along the bank of the lakes (as the aquatic birds are usually found around or in the lake) as followed by Gaston (1975) and Bibby *et al*. (2000). However, wherever necessary point count of birds was also made within the visible radius by stopping briefly for 2-3 min as followed by other workers (Froneman *et al*, 2001; Kaul and Howman, 1992; Urfi *et al*, 2005) and with help of binocular (Olympus 10x50). Identification, counting of the birds was made in the morning between 07:30 and 10:30 hr or in the afternoon between 15:00 and 18:00 hr, depending on the light conditions (Namgail *et al*, 2009).

Table 1: An annotated check list of aquatic avifauna of three lakes of Buldhana district (M.S.)

Scientific Name	Common Name	Status*	Occurrence	Food Habit*
Family: Podicipitidae				
<i>Tachybaptus ruficollis</i> (Pallas)	Little Grebe	R	C	O
Family: Phalacrocoracidae				
<i>Phalacrocorax niger</i> (Vieillot)	Little Cormorant	R	C	P
Family : Ardeidae				
<i>Ardea cinerea</i> (Linnaeus)	Grey Heron	RM	UC	P
<i>Ardeola grayii</i> (Sykes)	Indian Pond Heron	R	UC	P
<i>Bubulcus ibis</i> (Linnaeus)	Cattle Egret	R	C	P
<i>Mesophoyx intermedia</i> (Wagler)	Median Egret	R	C	P
<i>Egretta garzetta</i> (Linnaeus)	Little Egret	R	UC	P
Family: Ciconiidae				
<i>Mycteria leucocephala</i> (Pennat)	Painted Stork	RM	UC	C
<i>Anastomus oscitans</i> (Boddaert)	Asian Open-billed Stork	RM	UC	C
<i>Ciconia episcopus</i> (Boddaert)	White Necked Stork	R	C	C
Family: Threskornithidae				
<i>Threskornis melanocephalus</i> (Latham)	Oriental White Ibis	RM	R	O
Family: Anatidae				
<i>Anas poecilorhyncha</i> (J.R.Forester)	Spot Billed Duck	R	R	H
<i>Anas clypeata</i> (Linnaeus)	Northern Shoveller	M	R	C
Family: Rallidae				
<i>Amaurornis phoenicurus</i> (Pennat)	White Breasted Waterhen	R	R	O
<i>Gallinula chloropus</i> (Linnaeus)	Common Moorhen	R	R	O
<i>Porphyrio Porphyrio</i> (Linnaeus)	Purple Moorhen	R	R	O
Family: Recurvirostridae				
<i>Himantopus himantopus</i> (Linnaeus)	Black-winged Stilt	R	C	P
Family: Charidriidae				
<i>Vanellus malabaricus</i> (Boddaert)	Yellow-wattled lapwing	R	R	C
<i>Vanellus indicus</i> (Boddaert)	Red-wattled lapwing	R	R	C
Family: Scolopacidae				
<i>Tringa stagnatilis</i> (Linnaeus)	Marsh Sandpiper	M	UC	C
<i>Tringa glareola</i> (Linnaeus)	Wood Sandpiper	M	R	C
<i>Actitis hypoleucos</i> (Linnaeus)	Common Sandpiper	M	UC	C
Family: Laridae				
<i>Sterna aurantia</i> (J.E.Gray)	River Tern	RM	R	P
Family: Acledinidae				
<i>Acledo atthis</i> (Linnaeus)	Small Blue Kingfisher	R	UC	P
<i>Halcyon smyrnensis</i> (Linnaeus)	White-breasted Kingfisher	R	C	P
Family: Motacillidae				
<i>Anthus rufulus</i> (Vieillot)	Paddy field Pipit	R	C	I
<i>Motacilla Cinerea</i> (Tunstall)	Grey Wagtail	M	C	I
<i>Motacilla maderaspatensis</i> (Gmelin)	Large Pied Wagtail	R	C	I

*= Ali, S.,(2002)

Status- R- Resident; M- Migrant; RM- Resident Migrant;

Occurrence- C- Common; UC- Uncommon; R-Rare; NF- Not Found

Food Habit- C- Carnivorous; G- Granivorous; H- Herbivorous; I- Insectivorous; O- Omnivorous; P- Piscivorous

Surveys were conducted during 15 October to 15 November (2011). We observed and maintain the record of the birds of the area from different habitat including inhibited open areas, scrub of slopes, rim and lake basin. Birds observed in these habitats were segregated into Common(C), Uncommon (UC), Rare(R) and Not Found (NF).

The check list of species was prepared following Ali (2002), Manakadan and Pittie (2001) and Grimmett and Inskipp (2007).

RESULTS AND DISCUSSION

The studies on aquatic avifauna of Survey district revealed abundance of twenty-eight species of water birds of thirteen families (Table-1). As birds are highly mobile species of universe so it found equally distributed at all three lakes of Buldhana district. Of all the recorded families, Podicipitidae (1 Species), Phalacrocorax (1 species), Ardeidae (5 species), Ciconiidae (3 species), Threskornithidae (1 species), Anatidae (1 species), Rallidae (3 species), Recurvirostridae (1 species), Charadriidae (2 species) Scolopacidae (3 species), Laridae (1 species), Acledinidae (2 species) and Motacillidae (3 species) were recorded during 15 October to 15 November (2011).

From all the recorded birds, was residential birds of which 9 birds namely *Tachybaptus ruficollis*, *Phalacrocorax niger*, *Bubulcus ibis*, *Mesophoyx intermedia*, *Ciconia episcopus*, *Himantopus himantopus*, *Halcyon smyrnensis*, *Anthus rufulus*, *Motacilla maderaspatensis* shows common status. 3 birds namely *Ardeola grayii*, *Egretta garzetta*, *Acledo*

atthis was uncommon while 6 birds namely *Anas poecilorhyncha*, *Amauorornis phoenicurus*, *Gallinula chloropus*, *Porphyrio Porphyrio*, *Vanellus malabaricus*, *Vanellus indicus* were rare.

The resident migrant birds breed in one part of the area in one season and move to other parts within the state or country in a different season. 3 birds, *Ardea cinerea*, *Mycteria leucocephala*, *Anastomus oscitans* were uncommon while 2 birds namely *Sterna aurantia*, *Threskornis melanocephalusii* showed their rare status. Of all the recorded species, were migratory birds of which *Motacilla Cinerea* was common, *Tringa stagnatilis*, *Actitis hypoleucos* with uncommon status while *Anas clypeata*, *Tringa glareola* shows their rare status. The roosting birds like herons and egrets were also noted. The nearly same pattern of aquatic birds was also observed by Muhamed (2006).

From the above results it could be conclude that the abundance of avifauna indicates the healthy status of lakes owing the availability of water, safe habitat and food sources for both adults and nestlings and essential nesting/roosting sites in and around the lakes are important for the occurrence and abundance of aquatic bird populations. As water depth, quality and trophic structure are the important habitat characteristics that influence the abundance and diversity of aquatic birds in lakes, the proper and regular maintenance of these lakes would further increase the aquatic bird populations. The results of this study will help to conserve water bird populations in Buldhana district of Maharashtra, India.

LITERATURE CITED

- Ali S, 2002. *The book of Indian birds*. Oxford University Press, New Delhi, 13thEdn.
- Basavarajappa S, 2006. Avifauna of agro-ecosystems of maiden area of Karnataka. *Zoos' Print J.* **21**: 2217-2219.
- Bhatnagar SP, SK Shukla and M Bhatnagar, 2008. An annotated checklist of waterfowl of the Didwana Inland Saline Lake, Nagaur, Rajasthan, India. *News For Bird Watchers*, **48**: 53-55.
- Bibby CJ, DA Hill, ND Burgess and S Mustoe, 2000. *Bird census techniques*. Academic Press, London, 2nd Edn.
- Blair RB, 1999. Birds and butterflies along an urban gradient: Surrogate taxa for assessing biodiversity? *Ecol. Appl.*, **9**: 164-170.
- Custer TW and RG Osborne, 1977. Wading birds as biological indicators. Long Survey, US Fish and Wildlife service, Washington, DC.
- Emlen JT, 1974. An urban bird community in Tucson, Arizona: Derivation, structure, regulation. *The Condor*, **76**: 184-197.
- Froneman A, MJ Mangnall, RM Little and TM Crowe, 2001. Waterbird assemblages and associated habitat characteristics of farm ponds in the Western Cape, South Africa. *Biodiversity and Conservation*, **10**: 251-270.
- Gaston AJ, 1975. Methods for estimating bird populations. *J. Bombay Nat.Hist. Soc.*, **72**: 271-283.
- Gibbs JP, 1993. The importance of small wetlands for the persistence of local populations of wetland-associated animals. *Wetlands*, **13**: 25-31.
- Grimmett R and T Inskipp, 2007. *Birds of Southern India*. Om Books International, New Delhi, India.

- Hardy AR, PI Stanley and SPW Greeing, 1987.** Birds as indicator of the intensity of use of agricultural pesticide in UK. *In: The value of birds.* (Eds.: A.W. Diamond and F.N. Falion). Tech. Publ., **6**: 119-121.
- Joshua J and AJT Johnsingh, 1988.** Observations on birds on Mundanthurai Plateau, Tamil Nadu. *J. Bombay Nat. Hist. Soc.*, **85**: 565-577.
- Kaul R and S Howman, 1992.** Distribution and habitat requirements of thenorthern painted francolin *Fracolinus pictus pallidus* in Jamnagar, India. *Gibbier Faune Sauvage*, **9**: 483-492.
- Manakadan R and A Pittie, 2001.** Standardized common and scientific names of the birds of the Indian continent. *Buceros-Envis Newsl.*, **6**: 1-37.
- Muhamed JP, 2006.** A preliminary observation on the birds of Lonar crater, Buldhana district, maharashtra. *Zoo's Print J.*, **22(1)**: 2547-2550
- Namgail T, D Muddappa and TRS Raman, 2009.** Waterbird numbers at high altitude lakes in eastern Ladakh, India. *Wildfowl*, **59**: 137-144.
- Raghavaiah PS and P Davidar, 2007.** Status of migratory waterfowl of PulicatLake, Andhra Pradesh, India. *The Indian Forester*, **133**: 1312-1317.
- Rajshekhara S and MG Venkatesha, 2011.** Community of aquatic birds in lake of Bamglore, India , *J. of Environmental Biology*, **32(1)**: 77-83.
- Sahu HK and SD Rout, 2005.** Checklist of waterbirds in Mayurbhanj district, Orissa. *Zoos' Print J.*, **20**: 1992-1993.
- Surana R, BR Subba and KP Limbu, 2007.** Avian diversity during rehabilitation stage of Chimdi Lake, Sunsari, Nepal. *Our Nature*, **5**: 75-80.
- Urfi AJ, S Monalisa, A Kalam and Meganathan, 2005.** Counting birds in India: Methodologies and trends. *Curr. Sci.*, **89**: 1997-2003.